



ON THE COMB-TAILED *Neohahnia ernsti* (SIMON, 1898) (ARANEAE: HAHNIIDAE): REDESCRIPTION AND NEW DISTRIBUTION RECORDS

Nancy F. Lo-Man-Hung^{1*} , David F. Candiani²  and Alexandre B. Bonaldo³ 

¹ Programa de Pós-Graduação em Ciências Biológicas (Biologia Genética), Universidade de São Paulo, São Paulo, Brasil

² Metazoa Zoologia e Consultoria Científica LTDA, São Paulo, Brasil

³ Laboratório de Aracnologia, Coordenação de Zoologia, Museu Paraense Emílio Goeldi, Belém, Brasil

* Corresponding author: hahniidae@gmail.com

Fecha de recepción: 23 de octubre de 2024

Fecha de aceptación: 05 de diciembre de 2024

ABSTRACT

Neohahnia ernsti (Simon, 1898) is redescribed and illustrated based on newly examined material. For the first time, the male palp (retrolateral view) and female epigynum (ventral view) are documented. Scanning electron micrographs complement the species' description, and new distribution records are provided. *Neohahnia chalupas* Dupérré & Tapia, 2024 **syn. nov.** with *N. ernsti* (Simon, 1898) is proposed. Previously recorded only in its type locality, *N. ernsti* is now reported from several Neotropical regions, including location across the Caribbean, such as Saint Vincent Island, Puerto Rico, and Cuba, as well as Venezuela, Colombia, Ecuador, and Brazil. This study expands the known distribution of *N. ernsti* and offers morphological description into the species. Most specimens were collected from Neotropical rainforests, indicating a possible soil-dwelling habitat.

Keywords: dwarf sheet spiders; Hahniinae; taxonomy.

RESUMO

***Neohahnia ernsti* (Simon, 1898) (Araneae: Hahniidae): redescricao e novos registros de distribuição.** *Neohahnia ernsti* (Simon, 1898) é reescrita e ilustrada com base em material recentemente examinado. Pela primeira vez, o palpo masculino (visão retrolateral) e o epígino feminino (visão ventral) são documentados. Micrografias eletrônicas de varredura complementam a descrição da espécie, e novos registros de distribuição são fornecidos. *Neohahnia chalupas* Dupérré & Tapia, 2024 **syn. nov.** com *N. ernsti* (Simon, 1898) é proposto. Anteriormente registrada apenas para sua localidade tipo, *N. ernsti* agora é relatada em várias áreas neotropicais,

incluindo locais no Caribe, como Ilha de São Vicente, Porto Rico e Cuba, bem como Venezuela, Colômbia, Equador e Brasil. Este estudo expande a distribuição conhecida de *N. ernsti* e oferece descrição morfológica da espécie. A maioria dos espécimes foi coletada em florestas tropicais neotropicais, indicando um possível habitat de solo.

Palavras-chave: aranhas anãs de teia em lençol; Hahniinae; taxonomia.

INTRODUCTION

Neohahnia Mello-Leitão 1917 was originally described in the subfamily 'Hahniae' under Agelenidae, but it has been listed under Hahniidae since Gertsch (1934) and belonging to the subfamily Hahniinae as per Lehtinen (1967) until the present. Some authors have suggested a close relationship between *Neohahnia* and Neotropical species included in *Hahnia* (Mello-Leitão, 1917; Gertsch, 1934; Heimer and Müller, 1988), as well as *Alistra* and *Scotopsillus* (Mello-Leitão, 1917).

According to Mello-Leitão (1917), *Neohahnia* is closely related to *Alistra*, *Scotopsillus* and *Hahnia* because its anterior median eyes are smaller than the others. Additionally, *Neohahnia* is related to *Hahnia* due to the tracheal spiracle being closer to the spinnerets than the genital furrow (Schiapelli and Gerschman, 1958; Heimer and Müller, 1988). However, *Neohahnia* differs from all these genera because its anterior median eyes are contiguous and distinctly separated from anterior lateral eyes (Mello-Leitão, 1917).

Lehtinen (1967, p. 251) noted that the taxonomic status of this genus has remained somewhat obscure,



partly because the type specimen's preservation was unknown for him. Currently, *Neohahnia* Mello-Leitão 1917 comprises ten species (WSC, 2024). The type of specie *N. sylviae* was described based in juvenile exemplar from Rio de Janeiro, Brazil (Moreira et al., 2010). Mello-Leitão (1917) also described *N. palmicola*, which, like *N. sylviae*, was described based on a juvenile specimen from the same type locality. *N. chibcha* was described by Heimer and Müller (1988) based in male and female specimens from Colombia. Six species of *Neohahnia* were recently described from Ecuador (Dupérré and Tapia, 2024): *Neohahnia catleyi*, based on a female; *N. piemontana*, based on both male and female specimens; *N. pristirana*, based on both male and female specimens; *N. freibergeri*, based on both male and female specimens; *N. paramo*, based on both male and female specimens; and *N. chalupas*, based on a male.

Neohahnia ernsti (Simon, 1898) was described based in male and female specimens reported from St. Vincent, Venezuela. Later, Petrunkevitch (1929) described the structural characteristics for female species, as Simon had only provided details on the color pattern in his original descriptions. Petrunkevitch also expanded the know distribution of the species to Naguabo and Adjuntas in Puerto Rico. *N. ernsti* was originally placed in the genus *Hahnia* but was transferred to *Neohahnia* by Lehtinen (1967), who also provided figures of the male palp (ventral view) and female epigynum (dorsal view).

In the present paper, *N. ernsti* is recognized, redescribed, and illustrated. *Neohahnia chalupas* Dupérré and Tapia, 2024 is synonymized with *N. ernsti* (Simon, 1898). Male palp (retrolateral view) and female epigynum (ventral view) are illustrated for the first time. Scanning electron micrographs were taken as a complement to the species description, and new distribution records are provided.

MATERIAL AND METHODS

All morphological measurements, descriptions, drawings and color patterns were based on specimens immersed in 70% alcohol, using a stereomicroscope ZEISS SV11. The measurements were taken with an ocular micrometer and was used a conversion factor from the known measurement to convert the physical length of the marks on the scale in millimeters. Drawings were made with *camera lucida* attached on the stereomicroscope, male palp drawing were based on left pedipalp, female genitalia were first dissected following Levi (1965) then digested in trypsin enzyme (for digesting internal soft tissues) following modified protocol of Álvarez-Padilla & Hormiga (2007) and immersed in clove oil to be cleared for an easy examination of internal sclerotized structures. For scanning electron micrographs (SEM), the structures, after dissected, were cleaned for 3 min in alcohol in an

ultrasonic bath (Ultrasonic Cleaner USC 700) with water, then structures were submitted to dehydration by crescent alcohol concentration series (4 series: 80%, 85%, 90%, 96%) passed after to acetone 100% and then critical point dried from CO₂ using a critical point Dryer (Bal-Tec CPD 030 Critical Point Dryer) for 2h, the dried structures were fixed to an aluminum stub with a double-sided adhesive carbon tape, and sometimes glued in the tip of an entomological pins, the stubs were sputter coated with gold in a Bal-Tec SCD 005 Sputter Coater and placed in a vacuum desiccator until examination under a Philips XL 30 scanning electron microscope operated at 20kV. All the SEM protocol and images were made at Centro de Microscopia e Microanálises (CEMM) of the Pontifícia Universidade Católica do Rio Grande do Sul (PUCRS). Map produced with QGIS 3.40.3. Drawings and SEM images were processed with Open-Source image editing Gimp 2.6 (GNU Image Manipulation Program) and the final plates were edited with Open-Source vector graphics editor Inkscape 0.48.

Abbreviations used in the text and figures, as follows:

ALE Anterior lateral eyes; **ALS** Anterior lateral spinnerets; **AME** Anterior median eyes; **CD** Copulatory ducts; **CO** Copulatory openings; **cx** Coxae; **Cy** Cymbium; **d** Dorsal; **E** Embolus; **FD** Fertilization Ducts; **fe** Femur; **mt** Metatarsus; **p** Prolateral; **pa** Patella; **PA** Patellar apophysis; **PLE** Posterior lateral eyes; **PLS** Posterior lateral spinnerets; **PME** Posterior median eyes; **PMS** Posterior median spinnerets; **r** Retrolateral; **RTA** Retrolateral tibial apophysis; **T** Tegulum; **ta** Tarsus; **ti** Tibia.

Specimens examined are deposited in the following institutions (curators in parentheses):

IBSP Instituto Butantan, São Paulo, São Paulo (A.D. Brescovit)

MPEG Museu Paraense Emílio Goeldi, Belém, Pará (A.B. Bonaldo)

UFMG Universidade Federal de Minas Gerais, Belo Horizonte, Minas Gerais (A.J. Santos)

Taxonomy

Neohahnia ernsti (Simon, 1898)

Hahnia ernsti Simon 1898: 888 (male and female syntypes from Saint Vincent Island, in the Caribbean, also material from Caracas, Venezuela) (examined). WSC, 2024.

Hahnia ernsti Petrunkevitch 1929: 78-79, figs. 67-68 (four females from Puerto Rico: two from Naguabo, March 7, 1914; two from Adjuntas, June 8-13, 1915). Gertsch 1934: 9, 11 (two females from North America). Schiapelli and Gerschman de Pikelin 1958: 213 (cited from Central America). Opell and Beatty 1976: 421. Calvo 2000: 135-136 (one subadult female and one juvenile female)

from Cuba). *Lapsus calami*. WSC, 2024.

Neohahnia ernsti (Simon). Transference established by Lehtinen 1967: 251, figs. 366, 370. Heimer and Müller 1988: 229-230. *Lapsus calami*. WSC, 2024.

Neohahnia chalupas Dupérré and Tapia, 2024: 86-87 (male holotype from Ecuador, Napo Province, Tena, Colonso Chalupas Natural Reserve). WSC, 2024. **Syn. Nov.**

Type material. Syntypes of *Hahnia ernsti* Simon, 1898: male and female from St. Vincent Island and Caracas, Venezuela (MNHN 13284) (examined)

Note: ... "The tube labelled *B. antarctica* in Paris contains a single female of a *Neohahnia* sp., and it was interpreted as the holotype by Galiano" (in Schiapelli and Gerschman de Pikelin, 1959) (Lehtinen, 1967, p. 240-241). This information highlights historical taxonomic ambiguities in the group and the need to revisit and revise earlier type interpretations in the light of modern taxonomy, ensuring the accuracy of the redescrptions and distribution records.

Additional material examined. COLOMBIA: *Caqueta*: (Fazenda José Antonio Polo), 17.VI.2009, P. Lavelle et al., CSP214, 1♂ (MPEG 17760); BRAZIL: *Amazonas*: Coari (Base de Operações Geólogo Pedro de Moura, Porto Urucu), 4°52'31"S 65°10'27"W, 22.VIII.2008, S.C. Dias et al., Sid 1-206, 1♀ (MPEG 13455); Sid 1-204, 1♀ (MPEG 13456); *Pará*: L. Macambira, 94♂, 15♀ (MPEG 18429); Bragança, 22.VIII.2008, FST3-C3, 7♂, 1♀ (MPEG 13458); F5T1-C10, 7♂, 1♀ (MPEG 13462); F3T2-C9, 21♂, 1♀ (MPEG 13481); TT3-C9, 1♀ (MPEG 13485); TT3-C1, 1♀ (MPEG 13486); FST2-C14, 1♀ (MPEG 13490); FST1-C7, 9♂, 2♀ (MPEG 13491); FST3-C11, 3♂, 1♀ (MPEG 13493); FST3-C5, 3♂, 1♀ (MPEG 13494); FST3-C11, 2♀ (MPEG 13496); FST1-C17, 35♂, 1♀ (MPEG 13501); TT1-C5, 7♂, 1♀ (MPEG 13503); FST3-C17, 2♂ (MPEG 13515); FST2-C20, 6♂, 1♀ (MPEG 13517); FST2-C1, 6♂, 1♀ (MPEG 13518); FST2-C8, 3♂, 1♀ (MPEG 13523); FST2-C14, 1♀ (MPEG 13490); Belém (Ilha de Cotijuba), 01°14'S 48°35'W, 22.VIII.2008, Cotijuba 56, 1♀, 1j (MPEG 4717); Tailândia (Empresa Juruá Florestal. Fazenda Santa Marta), 21.I.2010, Equipe IPAN, Am 669, 1♀ (MPEG 17441); Am 225, 1♀ (MPEG 17442); Am 568, 1♀ (MPEG 17443); Am 574, 1♀ (MPEG 17445); Am 618, 1♀ (MPEG 17446); Am 656, 1♀ (MPEG 17447); Am 627, 1♀ (MPEG 17450); Am 705, 1♀ (MPEG 17452); Am 243, 1♀ (MPEG 17453); Am 245, 2♀ (MPEG 17454); Am 267, 1♀ (MPEG 17455); Am 276, 1♀ (MPEG 17457); Am 262, 2♀ (MPEG 17458); Am 606, 1♀ (MPEG 17460); Am 596, 1♀ (MPEG 17461); Am 264, 1♀ (MPEG 17462); Am 601, 1♀ (MPEG 17463); Am 637, 1♀ (MPEG 17465); Am 639, 1♀ (MPEG 17468); Am 699, 1♀ (MPEG 17470); Am 258, 2♀ (MPEG 17473); Am 247, 2♀ (MPEG 17475); Am 616, 1♀ (MPEG 17477); Am 701,

1♀ (MPEG 17478); Am 275, 1♀ (MPEG 17481); Am 713, 1♀ (MPEG 17482); Am 274, 2♀ (MPEG 17483); Am 578, 1♀ (MPEG 17484); Am 676, 1♀ (MPEG 17485); Am 610, 1♀ (MPEG 17486); Goianésia do Pará (Empresa Cikel Brasil Verde. Fazenda Rio Capim), 03°18'50"S 48°28'54" W, 21.I.2010, Equipe IPAN, Am 450, 1♀ (MPEG 17444); Am 76, 1♀ (MPEG 017449); Am 11, 1♀ (MPEG 17451); Am 439, 1♀ (MPEG 17459); Am 445, 1♀ (MPEG 17464); Am 9, 1♀ (MPEG 17467); Am 78, 1♀ (MPEG 17469); Am 407, 1♀ (MPEG 17472); Novo Repartimento (Empresa Juruá Florestal. Fazenda Arataú), 21.I.2010, Equipe IPAN, Am 483, 1♀ (MPEG 17448); Am 384, 1♀ (MPEG 17456); Am 532, 1♀ (MPEG 17466); Am 168, 1♀ (MPEG 17471); Am 510, 1♀ (MPEG 17474); Am 142, 1♀ (MPEG 17476); Am 372, 1♀ (MPEG 17479); Am 481, 1♀ (MPEG 17480); Am 345, 1♀ (MPEG 17487); Am 155, 1♀ (MPEG 17488); Am 359, 1♀ (MPEG 17489); Nova Ipixuna (Macaranduba), 682382.74/ 9469861.49, 17.VI.2009, P. Lavelle et al., BMB 1 DB4, 1♀ (MPEG 17759); *Tocantins*: Ananás, 06°13'34.70"S 52°25'2.39"W, U. Oliveira and M.D. Miranda, 2♂, 2♀ (UFMG 5751); Xambioá, 06°26'12.6"S 50°34'2.5"W, U. Oliveira and M.D. Miranda, 1♂ (UFMG 5701); Santa Fé do Araguaia, 06°43'41.6"S 48°48'8"W, U. Oliveira and M.D. Miranda, 1♂ (UFMG 5735); 1♂ (UFMG 5732); *Piauí*: José de Freitas (Fazenda Nazareth), 4°47'58.1"S 42°37'48.8"W, V.O. Costa, VOC(1)14, 1♀ (MPEG 9717); 08.VIII.2009, NZ TMSC 77, 1♂ (MPEG 17758); NZ TMSCO 168, 1♂ (MPEG 18177); NZ TMSCO 167, 1♂ (MPEG 18178); NZ TMSC 205, 2♂ (MPEG 18179); NZ TMSC 92, 1♀ (MPEG 18180); NZ TMSC 144, 1♂ (MPEG 18181); NZ TMSC 147, 1♂ (MPEG 18182); NZ TMSC 164, 1♂ (MPEG 18183); NZ TMSC 205, 1♂ (MPEG 18179); Maranhão: Caxias (APA do Inhamum), 2017-2018, I. Lima, Am 475 pitfall, 1♀ (IBUSP 246475); Bahia: Ilhéus (Campus UFSC, Cabruaca), 20.VII.1999, M.F. Dias, 1♀ (IBUSP 24226); *Mato Grosso*: Cotriguaçu (Ilhas do Rio Juruá), 09°52'40"S 58°12'38"W, A.J. Santos, 1♂, 3♀ (UFMG 7859); (Fazenda São Nicolau), 09°50'24"S 58°14'54"W, A.J. Santos, 1♂, 2♀, 1j (UFMG 7708); 1♂ (UFMG 7709); Canarana/Querência (Fazenda Tanguro, antiga Fazenda Morro Azul), 13°04'25.1"S 52°22'54.9"W, 22.VIII.2008, D.F. Candiani and N.F. Lo-Man-Hung, Tang 416, 1♀ (MPEG 13452); Nossa Senhora do Livramento (Distrito de Pirizal, Fazenda Retiro Novo), 16°22'1"S 56°17'58"W, F.S.F. Leite, 1♀ (UFMG 8771); *Mato Grosso do Sul*: Três Lagoas (Horto Barra do Moeda), 20°57'0"S 51°47'0"W, M. Uehara-Prado, 1♂, 4♀ (UFMG 5684); 13♂, 5♀ (UFMG 5685); 1♂, 1♀ (UFMG 5686); 6♂, 3♀ (UFMG 5687); 3♂, 4♀ (UFMG 5688); 6♀ (UFMG 5689); 11♂, 2♀, 1j (UFMG 5690); Brasilândia (Horto Rio Verde), 20°50'S 51°40'W, M. Uehara-Prado, 3♂, 2♀ (UFMG 5713); (Usina Hidrelétrica Sérgio Motta), 31.VII-07.VIII.2000, Equipe IBSP, Etapa II, pitfall 03, 2♀ (IBSP 30826); Etapa II, pitfall 43, 2♂ (IBSP 30841); Etapa II, pitfall 19, 2♂ (IBSP 30854); Etapa II, pitfall 43, 1♂ (IBSP 30863); Etapa II,



pitfall 46, 1♂ (IBSP 30864); (Usina Hidrelétrica Sérgio Motta, Fazenda Cizalpira), 15-9.VIII.2000, C. Fukushima and J.P. Guadanucci, Etapa II, pitfall 37, 1♂ (IBSP 35376); Etapa II, pitfall 23, 1♂ (IBSP 35378); Etapa II, pitfall 25, 1♂ (IBSP 35393); MS 21, 2♂ (IBSP 35394); Etapa II, pitfall 6, 1♂, 2♀ (IBSP 35518); 2000, Equipe IBSP, pitfall 25, MS 16, 1♀ (IBSP 30454); pitfall 2, MS 26 1♀ (IBSP 30456); MS 24, 1♂, 1♀ (IBSP 30471); pitfall 2, 1♀ (IBSP 35519); MS 24, 1♂ (IBSP 35527); MS 8, 1♂ (IBSP 35536); pitfall 47, 1♂, 1♀ (IBSP 35373); MS 41, 3♂ (IBSP 35374); MS 51, 4♂ (IBSP 35375); MS 50, 2♂ (IBSP 35389); MS 35, 4♂ (IBSP 35408); MS 38, 1♂ (IBSP 35410); MS 05, 3♂ (IBSP 35532); *Minas Gerais*: Jaboticatubas (Parque Nacional da Serra do Cipó), 19°15'S 43°31'W; 1♂ (IBSP 159025); 1♂ (IBSP 159026); 1♂ (IBSP 159027); 1♂ (IBSP 159028); 1♂ (IBSP 159029); 1♂ (IBSP 159030); 1♂ (IBSP 159031); 1♂ (IBSP 159032); 1♂ (IBSP 159033); 1♂ (IBSP 159034); 1♂ (IBSP 159035); 1♂ (IBSP 159036); 1♂ (IBSP 159037); 1♂ (IBSP 159038); 1♂ (IBSP 159039); 1♂ (IBSP 159040); 1♂ (IBSP 159041); 2♂ (IBSP 159042); 1♂ (IBSP 159043); 1♂ (IBSP 159044); 1♂ (IBSP 159045); 1♂ (IBSP 159046); 1♂ (IBSP 159047); 1♂ (IBSP 159048); 1♂ (IBSP 159050); 1♂ (IBSP 159051); 1♂ (IBSP 159052); 1♂ (IBSP 159053); 1♂ (IBSP 159055); 1♂ (IBSP 159057); 1♀ (IBSP 159058); 1♂ (IBSP 159059); 1♂ (IBSP 159060); 1♂ (IBSP 159061); 1♂ (IBSP 159062); 1♂ (IBSP 159063); 2♂ (IBSP 159064); 1♂ (IBSP 159065); 1♂ (IBSP 159066); 2♂ (IBSP 159067); 1♂ (IBSP 159069); 1♂ (IBSP 159070); 07-14.X.2002, coleta 3: Am 100, 2♂ (IBSP 159123); Am 37, 1♂ (IBSP 159136); Am 73, 1♂ (IBSP 159183); 03-09/I.2003, equipe Biota, Am 63 pitfall: campo rupestre, coleta 4, 2♂ (IBSP 159184); Am 49, 2♂ (IBSP 159185); Am 55 pitfall, 4♂ (IBSP 159186); Am 50, 1♂ (IBSP 159188); Am 33, 1♂ (IBSP 159190); Am 150, 1♂ (IBSP 159192); Am 36, 2♂ (IBSP 159193); Am 15 pitfall: campo rupestre, 1♂ (IBSP 159194); Am 71, 2♂ (IBSP 159196); Am 42, 1♂ (IBSP 159197); Am 48, 1♂ (IBSP 159198); Am 47, 1♂ (IBSP 159199); Am 62, 1♂ (IBSP 159200); Am 116, 1♂ (IBSP 159203); Am 59, 1♂ (IBSP 159205); Am 112, 1♂ (IBSP 159206); Am 62, 1♂ (IBSP 159207); Am 53, 1♂ (IBSP 159208); Am 94, 2♂ (IBSP 159209); Am 120, 1♂ (IBSP 159211); Am 70, 3♂ (IBSP 159212); Am 96 pitfall, 1♂ (IBSP 159214); Am 137, 2♂ (IBSP 159215); Am 99, 1♂ (IBSP 159216); Am 75, 2♂ (IBSP 159217); Am 58, 1♂ (IBSP 159219); Am 9 pitfall: campo rupestre, coleta 4, 1♂ (IBSP 159221); Am 5, 1♂ (IBSP 159223); Am 138, 1♂ (IBSP 159225); Am 29, 1♂ (IBSP 159229); Am 93 pitfall: campo rupestre, coleta 4, 1♂ (IBSP 159226); Am 87, 2♂ (IBSP 159227); Am 145 pitfall: campo rupestre, coleta 4, 1♂ (IBSP 159228); Am 148, 1♂ (IBSP 159233); 2♂ (IBSP 159235); Am 142, 1♂ (IBSP 159237); Am 80, 1♂ (IBSP 159238); Am 83, 2♂ (IBSP 159239); Am 126 pitfall, 1♂ (IBSP 159241);

Am 119, 1♂ (IBSP 159242); Am 108, 1♂ (IBSP 159245); Am 105, 1♂ (IBSP 159246); Am 19 pitfall: campo rupestre, coleta 4, 2♂ (IBSP 159247); 1♂ (IBSP 159249); Am 24, 1♂ (IBSP 159924); Santana do Riacho (Parque Nacional da Serra do Cipó, Portaria Palácio), 19°15'S 43°31'W, E.S.S. Álvares and E.O. Machado, 1♀ (UFMG 1413); Belo Horizonte (Estação Ecológica da UFMG), 19°52'28"S 43°58'22"W, E.S.S. Álvares and E.O. Machado, 6♂ (UFMG 1587); E.S.S. Álvares and E.O. Machado, 3♂, 7♀ (UFMG 1588); E.S.S. Álvares, 3♂, 1♀ (UFMG 5016); E.S.S. Álvares et al., 1♂ (UFMG 5436); E.S.S. Álvares, 4♂ (UFMG 6030); E.S.S. Álvares, 3♂ (UFMG 6049); I.L.E. Magalhães, 2♂ (UFMG 8528); Ouro Preto (Parque Estadual do Itacolomi), 20°22'S 32°32'W, K.P. Santos et al., 1♀ (UFMG 2022); 1♂ (UFMG 2025); 1♀ (UFMG 2105); 1♂, 2♀ (UFMG 2284); 62♂ (UFMG 2332); 56♂, 37♀ (UFMG 2333); 4♂ (UFMG 2334); (Estação Ecológica de Tripuí), Am P1-6, 3♂, 3♀ (IBSP 159020); Am P8-6, 3♂ (IBSP 159021); *São Paulo*: São Paulo (Parque da Previdência), 16-23.XI.1999, D.F. Candiani, Verão PREV IV-37, 1♀ (IBSP 42105); Verão PREV IV-43, 1♀ (IBSP 42106); Verão PREV IV-35, 1♂ (IBSP 42107); Verão PREV IV-24, 1♂, 1♀ (IBSP 42108); Verão PREV IV-34, 1♀ (IBSP 42109); Verão PREV IV-22, 1♂ (IBSP 42112); Verão PREV IV-12, 1♀ (IBSP 42115); Inverno PREV II-12, 1♂ (IBSP 42120); Inverno PREV II-21, 1♂ (IBSP 42121); Inverno PREV II-47, 1♂ (IBSP 42126); Inverno PREV II-43, 1♂ (IBSP 42127); Inverno PREV II-35, 1♂ (IBSP 42129); Primavera PREV III-40, 1♂ (IBSP 42135); Primavera PREV III-02, 1♂, 1♀ (IBSP 42142); V.2000-II.2001, pitfall 1♀ (IBSP 76547); Santa Rita do Passo Quatro (Parque Estadual de Vassununga), 21°43'S 47°35'W, 19-24.III.2002, Equipe Biota, 2♀ (IBSP 60121); Campos do Jordão (Parque Estadual de Campos do Jordão), 22°43'S 45°27'W, alt 1557 m, IX.2004 – IX.2005, D. Baretta, Am 10, 1♂ (IBSP 88022); Am 65, 1♂ (IBSP 88023); Am 65, 1♂ (IBSP 88035); *Rio de Janeiro*: Petrópolis (Fazenda Ranchinho da Roça), 15-16.VIII.2001, Equipe Biota, Am 14 cascata pitfall traps, 1♂ (IBSP 157041); Am 134 cascata pitfall traps, 1♂ (IBSP 157042); Am 35 PF cascata pitfall traps, 3♂, 1♀ (IBSP 157045); Am 21 cascata pitfall traps, 1♂ (IBSP 157051); Am 01 cascata pitfall traps, 1♂ (IBSP 157053); Am 9 PF cascata pitfall traps, 1♂, 1♀ (IBSP 157055); Am 02 cascata pitfall traps, 2♂ (IBSP 157063); Am 33 cascata pitfall traps, 1♀ (IBSP 157065); Am 38 cascata pitfall traps, 2♂, 1♀ (IBSP 157066); (Fazenda Ranchinho, Ponto da Roça Km 78,5 – BR 040), 2001, F.S. Cunha, 1♀ (IBSP 35884); *Paraná*: Cornélio Procopio (PE Mata São Francisco), 28°08'47"S 50°34'19"W, V.2009-IV.2010, N. Cipola, Am:133, 1♀ (IBSP 163861); Am:180, 1♀ (IBSP 163862); Am:47, 1♂ (IBSP 164673); Am:89, 4♂ (IBSP 164674); Am:69, 4♂, 1♀ (IBSP 164675); Am:88 1♂ (IBSP 164676); Am:26, 2♂ (IBSP 164677); Am:68, 2♂ (IBSP 164678); Am:158, 1♂ (IBSP 164679); Am:48, 1♂ (IBSP 164680); Am:182,



1♂ (IBSP 164681); Rio Grande (Fazenda Experimental Grelha Azul), 25°40'16"S 49°16'44"W, IX.2010-X.2011, J. Ricetti, FEGA 183, 1♀ (IBUSP 220616); IRATI 270, 1♀ (IBUSP 220618);

Diagnosis. *Neohahnia ernsti* shares several characteristics with other species, particularly *N. chibcha*, *N. piemontana*, *N. pristirana*, *N. freibergi*, and *N. paramo*, but can be distinguished by a combination of key features. The males of *N. ernsti*, like those of *N. chibcha*, *N. piemontana*, *N. pristirana*, have a rounded cymbium. Additionally, similar to *N. chibcha*, *N. ernsti* males have an embolus that curves in a clockwise direction around the bulbus in a complete circle (Fig. 2), lying in a furrow on the tegulum, without a median apophysis, and exhibiting a long, curved retrolateral tibial apophysis (RTA). However, *N. ernsti* males differ from other species in the direction of the RTA and the shape of the embolus. In *N. chibcha*, the embolus curves in an S-shape toward the cymbium (Heimer and Müller, 1988: fig. 4), while in *N. piemontana*, the RTA is larger (Dupérré and Tapia, 2024: fig. 11B), and in *N. pristirana*, the RTA is shorter and more angular than in *N. ernsti*, with a different orientation (Dupérré & Tapia, 2024: fig. 18B). In *N. chibcha*, the RTA points toward a curved patellar apophysis (PA) (Heimer and Müller, 1988: fig. 3), in *N. piemontana*, the PA is bidentate (Dupérré & Tapia, 2024: fig. 11B), and in *N. freibergi*, the PA is single-pointed (Dupérré and Tapia, 2024: fig. 20B). In contrast, in *N. ernsti* the embolus curve in a slight V-shape in the cymbial furrow, between the bulbus and the cymbium laterally, and the RTA is thin and semicircular, direct against the PA which is also curved with 1-3 hook (Figs. 3-4). Females of *N. ernsti* resemble females of *N. chibcha* in having a simple epigynum, but they differ in the shape and direction of the copulatory openings, copulatory ducts, and fertilization tubes. In *N. chibcha*, the copulatory openings are rectangular (Heimer & Müller, 1988: fig. 5), and the copulatory ducts form highly coiled, irregular loops, which terminate in connection with the posterior spermathecae. The fertilization tubes are small and lateral (Heimer and Müller, 1988: fig. 6). Whereas females of *N. ernsti* present a simple epigynum, with paired copulatory openings (or "pockets") that are very conspicuous, and the copulatory ducts are almost globular with small central fertilization tubes (Figs. 5-6). When compared to *N. piemontana*, *N. ernsti* females can be distinguished by their copulatory openings, which are less separated (Fig. 5), whereas in *N. piemontana*, the openings are more widely spaced (Dupérré and Tapia, 2024: fig. 11D). Females of *N. pristirana* have large, rounded secondary spermathecae and bean-shaped primary spermathecae (Dupérré & Tapia, 2024: fig. 18D), while *N. freibergi* has elongated, joined copulatory openings accompanied by a small triangular hood (Dupérré and Tapia, 2024: fig. 20C, D). In contrast, *N. ernsti* females have copulatory ducts more globular. Furthermore,

compared to *N. paramo*, the internal genitalia and spermathecae of *N. ernsti* differ in the arrangement of the copulatory ducts and the absence of more defined loops (Dupérré and Tapia, 2024: fig. 22D), which is observed in *N. paramo*.

Description. Male (MPEG 18179). Total length 1.64, Cephalothorax 0.70 long, 0.60 wide. Eye diameters and interdistances: AME 0.08, ALE 0.06, PME 0.06, PLE 0.04, AME-AME 0 (contiguous), AME-ALE 0.01, PME-PME 0.06, PME-PLE 0.02, ALE-PLE 0 (contiguous). Clypeus shallow 0.08. Spinnerets total length and interdistances: ALS 0.24, PLS 0.36, MLS 0.14, ALS-PLS 0.20, ALS-MLS 0.04, MLS-MLS 0.02. Legs length: Coxae I 0.22, II 0.22, III 0.22, IV 0.24, Trochanter I 0.08, II 0.08, III 0.08, IV 0.10, Femur I 0.56, II 0.56, III 0.50, IV 0.60, Patella I 0.20, II 0.20, III 0.20, IV 0.20, Tibia I 0.50, II 0.44, III 0.40, IV 0.46, Metatarsus I 0.38, II 0.38, III 0.38, IV 0.50, Tarsus I 0.32, II 0.34, III 0.36, IV 0.42. Dorsal carapace pale brown with prominent border mottled dark brown, dark brown bands in *pars thoracica*. *Pars cephalica* a little darker than *pars thoracica* (see Fig. 1 for mottled pattern). Carapace with plate structure (Fig. 7-8). Longitudinal black thoracic low groove (Fig. 8). Eyes bordered in black, except PME. Carapace almost smooth, with a few long setae in a longitudinal row between fovea and PME, some long setae scattered in eyes (Fig. 7). Chelicerae pale brown mottled dark brown, with two promarginal (Fig. 9, 16) and one retromarginal teeth, without stridulatory files. Endites rectangle shape (Figs. 10-11), 0.20 long, 0.16 wide, with same color's pattern of chelicerae, long setae scattered, serrula in a single row of teeth (Figs. 12-13, 18, 20). Labium trapezoidal shape (Fig. 14), 0.04 long, 0.08 wide, pale brown with few long setae. Sternum circular shape, 0.42 long, 0.42 wide, pale brown with soft border dark brown, long setae in border and short setae scattered in the middle of sternum. Coxae and trochanters pale beige (especially in ventral view), remaining segments of legs and pedipalps brown-orange, setae, macrosetae and trichobothria, three claws. Dorsal abdomen dark brown with large and well-defined beige chevrons in the middle of abdomen and pale beige points around it (see Fig. 1 for chevrons pattern). Ventral abdomen unicolor, pale, same color as sternum, without chevrons or points, few setae (about eight) in epigastric furrow and book lungs base, tracheal spiracle in the middle of abdomen, same distance of spinnerets base as epigastric furrow. Spinnerets with modified topology, arranged in transversal row (Fig. 23), PMS, ALS and MLS with same color as ventral abdomen, pale brown, PMS and ALS segmented (Fig. 23), PMS longer than others (Fig. 23), gland spigots PLS cylindrical spigots and piriform spigots (Figs. 24-28, 35), ALS gland spigots; cylindrical spigots and piriform spigots (Figs. 29-30, 36-37), MLS gland spigots; cylindrical spigots and piriform spigots (Figs. 31-34, 38). Cymbium rounded, embolus thin and long (Fig. 2), without





Fig. 1-6. *Neohahnia ernsti* (Simon 1898). 1, Habitus male, dorsal view; 2, Male palp ventral view; 3, Male palp retrolateral view; 4, Male tibia and patella palp; 5, Female epigynum dorsal view; 6, Female epigynum ventral view.


median apophysis, shape of retrolateral tibial apophysis, thin, long and curved against the patellar apophysis which is also curved (Fig. 3-4). Retroventral tibial apophysis.


Description. Female (MPEG 13490). Total length 1.86, Cephalothorax 0.66 long, 0.56 wide. Eye diameters and interdistances: AME 0.06, ALE 0.06, PME 0.06, PLE 0.04, AME-AME 0.02, AME-ALE 0.01, PME-PME 0.06, PME-PLE 0.02, ALE-PLE 0 (contiguous). Spinnerets total length and interdistances: ALS 0.22, PLS 0.42, MLS 0.14, ALS-PLS 0.20, ALS-MLS 0.06, MLS-MLS 0.04. Legs length: Coxae I 0.22, II 0.20, III 0.20, IV 0.22, Trochanter I 0.10, II 0.08, III 0.08, IV 0.10, Femur I 0.56, II 0.54, III 0.48, IV 0.60, Patella I 0.22, II 0.20, III 0.18, IV 0.20, Tibia I 0.46, II 0.40, III 0.38, IV 0.58, Metatarsus I 0.38, II 0.32, III 0.34, IV 0.50, Tarsus I 0.32, II 0.32, III 0.38, IV 0.42.

Color and description same as male, except dorsal carapace with border mottled dark brown and dark brown bands in *pars thoracica*, not too defined as in male. Endites (Figs. 15, 17, 19) 0.12 long, 0.10 wide. Labium 0.06 long, 0.10 wide (Figs. 15, 21). Sternum ovoid shape (Fig. 22), 0.32 long, 0.40 wide. Coxae and trochanters pale brown (especially in ventral view), remaining segments of legs and pedipalps with same color pattern of carapace, pale brown with mottled dark brown. Dorsal abdomen chevrons and points not too defined as in male. Epigynum larger (almost the same length of book lung, almost twice wider than book lung), globose (Fig. 5-6), copulatory opening next to pedicel.

Variation. Females: size, color pale, mottled pattern not too defined; Males: size, color not too brown-orange.








Natural history. Most individuals were collected in Neotropical rain forests. When collected in Amazon, the places were characterized by tall dense jungle on dry soil. Most individuals were collected by Winkler's extractor, pitfall traps and Tropical Soil Biology and Fertility Program (TSBF) method, suggesting the specimens has a soil habitat.

Distribution. Island in Saint Vincent and the Grenadines (type locality) (Simon, 1898), Venezuela, (Simon, 1898), Puerto Rico: Naguaboi, Adjuntas (Petrunkevitch 1929), Cuba (Calvo 2000), Colombia, Ecuador (Dupérré & Tapia, 2024), Brazil: Pará, Tocantins, Piauí, Maranhão, Bahia, Mato Grosso, Mato Grosso do Sul, Minas Gerais, São Paulo, Rio de Janeiro, Paraná (Map 1).

ACKNOWLEDGMENTS

Thanks to Nadine Dupérré for the information on *Neohahnia chalupas* Dupérré & Tapia, 2024. Special thanks to the Centro de Microscopia e Microanálises (CEMM) of Pontifícia Universidade Católica do Rio Grande do Sul (PUCRS) for their support. We also extend our gratitude to João Lucas Chavari for providing records from Paraná and to Rafael Francisco for examination the type specimen. This research was supported by the Brazilian Council for the Development of Science (CNPq), through the M.Sc. scholarship process 130775/2011-8 awarded to NFLMH and PQ-grant #307165/2022-1 to ABB.





REFERENCES

- Álvarez-Padilla, F., & Hormiga, G. (2007). A protocol for digesting soft tissues and mounting spiders for scanning electron microscopy. *Journal of Arachnology*, 35, 538-542.
- Calvo, A.F.A. (2000). Primer registro de la familia Hahniidae (Arachnida: Araneae) para Cuba. *Avicennia*, 12/13, 135-136.
- Dupérré, N., & Tapia, E. (2024). First record of the family Hahniidae in Ecuador with description of thirteen new species and three new genera (Araneae: Hahniidae). *Taxonomy*, 4(1), 53-111.
- Gertsch, W.J. (1934). Some American spiders of the family Hahniidae. *American Museum Novitates*, 712, 1-32.
- Heimer, S., & Müller, H.G. (1988). Spiders from Colombia VIII. *Neohahnia chibcha* n. sp. from the Sierra Nevada mountains, northern Colombia (Arachnida, Araneida, Hahniidae). *Bulletin of the British Arachnological Society*, 7, 229-230.
- Lehtinen, P.T. (1967). Classification of the cribellate spiders and some allied families, with notes on the evolution of Araneomorphae. *Annales Zoologici Fennici*, 4(3), 1-199.
- Levi, H.W. (1965). Techniques for the study of spider genitalia. *Psyche*, 72, 152-158.
- Mello-Leitão, C.F. 1917. Gêneros e espécies novas de araneidos. *Archivos da Escola Superior de Agricultura e Medicina Veterinária*, 1, 3-19.
- Moreira, T.S., Baptista, R.L.C., Kury A.B., Giupponi, A.P.L., Buckup, E.H., & Brescovit A.D. (2010). Annotated check list of Arachnidae type specimens deposited in the Museu Nacional, Rio de Janeiro. II – Araneae. *Zootaxa*, 2588, 1-91.
- Opell, B.D., & J.A. (1976). Beatty. The Nearctic Hahniidae (Arachnida: Araneae). *Bulletin of the Museum of Comparative Zoology at Harvard*, 147, 393-433.
- QGIS.org, 2025. QGIS Geographic Information System. QGIS Association. <http://www.qgis.org>
- Petrunkévitch, A. (1929). The spiders of Porto Rico. Part one. *Transactions of the Connecticut Academy of Arts and Sciences*, 30, 1-158.
- Schiapelli, R.D., & B.S. Gerschman de P. (1958). Arañas argentinas III. Arañas de Misiones. *Revista del Museo Argentino de Ciencias Naturales "Bernardino Rivadavia"*, 3, 187-231.

- Schiapelli, R.D., & B.S. Gerschman de P. (1959). Aclaración sobre los caracteres genericos del género *Bigois* Simon, 1898 (Araneae, Hahniidae). *Revista de La Sociedad Entomológica Argentina*, XXI, 129-132.
- Simon, E. (1898). *On the spiders of the Island of St. Vincent Part III*. Proceedings of the Zoological Society of London 860-890.
- World Spider Catalog. 2024. *World Spider Catalog*. Natural History Museum Bern, online at <http://wsc.nmbe.ch>, version 25.5, accessed in October 2024. doi: 10.24436/2

Section Editors:
Anita Aisenberg, Macarena González,
Carolina Rojas-Buffer